arXiv:cond-mat/0110222v1 11 Oct 2001

Shake-up and shake-off excitations with associated electron losses in X-ray studies of proteins

Petter Persson [1], Sten Lunell [1], Abraham Szöke [2,3], Beata Ziaja [2,4,5] and Janos Hajdu [2,6]

[1] *Department of Quantum Chemistry, Uppsala University, Box 518, S-751 20 Uppsala, Sweden*

[2] *Department of Biochemistry, Biomedical Centre, Box 576, Uppsala University, S-75123 Uppsala, Sweden*

[3] *Lawrence Livermore National Laboratory, Livermore, CA 94551, USA*

[4] *Department of Theoretical Physics, Institute of Nuclear Physics, Radzikowskiego 152, 31-342 Cracow, Poland*

[5] *High Energy Physics, Uppsala University, P.O. Box 535, S-75121 Uppsala, Sweden*

Corresponding author:
Janos Hajdu, Department of Biochemistry, Biomedical Centre,
Box 576, Uppsala University, S-75123 Uppsala, Sweden
Tel:+4618 4714999, Fax:+4618 511755, E-mail:hajdu@xray.bmc.uu.se

Manuscript information:
Number of pages (double-sided version with tables and figure legend on separate pages): 19 , Number of figures:1, Number of tables:2

Submission information:
Diskette of the manuscript included, manuscript written in LaTeX, ASCII file extracted; hard copy double-spaced

[6] e -mail:petter.persson@kvac.uu.se, sten.lunell@kvac.uu.se, szoke1@llnl.gov, ziaja@tsl.uu.se, hajdu@xray.bmc.uu.se

Abstract

Photoionisation of an atom by X-rays usually removes an inner-shell electron from the atom, leaving behind a perturbed "hollow ion" whose relaxation may take different routes. In light elements, emission of an Auger electron is common. However, the energy and the total number of electrons released from the atom may be modulated by shake-up and shake-off effects. When the inner-shell electron leaves, the outer-shell electrons may find themselves in a state that is not an eigen-state of the atom in its surroundings. The resulting collective excitation is called shake-up. If this process also involves the release of low energy electrons from the outer shell, then the process is called shake-off. It is not clear how significant shake-up and shake-off contributions are to the overall ionisation of biological materials like proteins. In particular, the interaction between the out-going electron and the remaining system depends on the chemical environment of the atom, which can be studied by quantum chemical methods. Here we present calculations on model compounds to represent the most common chemical environments in proteins. The results show that the shake-up and shake-off processes affect about 20% of all emissions from nitrogen, 30% from carbon, 40% from oxygen, and 23% from sulphur. Triple and higher ionisations are rare for carbon, nitrogen and oxygen, but are frequent for sulphur. The findings are relevant to the design of biological experiments at emerging X-ray free-electron lasers.

Keywords: X-rays, photoionisation, shake-up, shake-off, Auger emission, radiation damage, peptides, proteins

Computer simulations (Neutze *et al.*, 2000) show that ultrashort and high-intensity X-ray pulses, as those expected from presently developed free-electron lasers (Winick, 1995; Wiik, 1997), may provide structural information from large protein molecules and assemblies before radiation damage destroys them. Estimation of radiation damage as a function of X-ray photon energy, pulse length, integrated pulse intensity and sample size was obtained in the framework of a radiation damage model, where the effects of atom-photon and ion-ion interaction were taken into account. Photons of 1 Å wavelength, corresponding to a photon energy of about 12 keV, interact with atoms mainly via the photoelectric effect (Dyson, 1973) (for carbon the photoelectric cross-section is ~ 10 times higher than the corresponding elastic cross-section at this wavelength (Hubbel *et al.*, 1980)), and thus the photoelectric effect is the main source of radiation damage with X-rays.

Photoionisation may proceed either through the ejection of an outer-shell electron or through the ejection of an inner-shell electron (Dyson, 1973). Outer-shell photo-events eject a single electron from the atom with an energy equivalent to the energy of the incoming photon minus the shell binding energy and the recoil energy. With photon energies of around 12 keV (or about 1 Å wavelength), outer-shell events are rare in carbon, nitrogen, oxygen and sulphur, and represent less than 5% of all photoionisation events.

A more frequent type of photoionisation with X-rays involves the ejection of an inner-shell electron from the atom (around 95-97% of photoionisations remove a K-shell electron from carbon, nitrogen, oxygen and sulphur). Typical K-hole life times are of the order $1-10$ fs (Krause & Oliver, 1979), and the hollow ion relaxes through an electron falling from a higher shell into the vacant hole. In light elements, the energy of this electron is given to another electron which is then also ejected from the atom through the Auger effect. Core ionisation, however, constitutes a strong perturbation of the molecule, and may be accompanied by significant electronic effects (Siegbahn *et al.*, 1969). Firstly, the valence electrons often relax significantly to compensate for the presence of the positive core hole. The departing photoelectron can interact with these relaxing electrons, and lose kinetic energy in the process, thus forcing the system into an excited state (a process called *shake-up*). In some cases, the excitation may result in the ejection of low energy outer-shell electrons from the atom (a process called *shake-off*). The multiple excitation lifetimes for light elements are comparable to core-hole lifetimes. The interaction between the out-going electron and the remaining system depends on the chemical environment of the atom, and a description of these interactions requires quantum chemical calculations. The following key processes need to be taken into account:

(i) PHOTOEMISSION FOLLOWED BY SINGLE AUGER DECAY.
Ejected K-shell electrons with the highest possible kinetic energy correspond to a situation where the remaining system is left in the ground state, i.e. the difference between the incoming photon energy and the energy of the ejected photoelectron equals the chemical binding energy and the recoil energy. In light elements like carbon, nitrogen, oxygen and sulphur, such clean photoemissions are followed by Auger emission, and the atom becomes doubly ionised (Krause & Oliver, 1979). In heavier elements X-ray fluorescence dominates.

(ii) SHAKE-RELATED PROCESSES. When photoionisation ejects an inner-shell electron so fast that the outer-shell electrons have no time to relax, the situation gets similar to beta decay, whereby the nuclear charge suddenly increases by one unit. Quantum mechanics describes such a state as a superposition of proper eigen-states, that include states where one or more of the electrons may be unbound. Interactions between the departing photoelectron and the electrons left behind may reduce the kinetic energy of the photoelectron, and deposit energy into the system. The perturbation is called shake-up if they refer to an excitation in the final system, or shake-off if the result is the loss of one or more outer shell electrons from the ion. The relative contributions from these two processes have been found to be comparable in noble gases (Svensson *et al.*, 1988; Armen *et al.*, 1985; Wark *et al.*, 1991). The following shake-related phenomena need to be considered:

(a) Photoemission accompanied by shake-up excitation with Auger decay. This process reduces the energy of the photoelectron slightly (about 10-40 eV), and the energy difference can either be absorbed by the atom or added to the energy of the Auger electron. At the end, the atom becomes doubly ionised.

(b) Photoemission accompanied by a shake-off event. In the vicinity of a hole, a vacancy and a free electron are created. The shake electron has around 10-100 eV energy, and the atom becomes doubly ionised.

(c) Photoemission accompanied by double Auger decay, which may proceed via different mechanisms, and may result in the triple ionisation of the atom. These mechanisms include Auger cascading, single Auger decay combined with shake-off emission, and virtual inelastic scattering (for a full description see Amusia *et al.*, 1992). In this case, the energies of the two ejected electrons are asymmetrically distributed. For instance, for Ne (transition $1s^{-1} \rightarrow 2s^{-2}2p^{-1} + q_1 + q_2$) the total available kinetic energy is about 650 eV, and the most probable case is shaking off a slow electron and Auger emission of a fast electron with $E_{shake-off} \ll E_{Auger}$ (Amusia *et al.*, 1992).

In the initial damage model of proteins (Neutze *et al.*, 2000) only outer-shell pho-

toionisation and inner-shell photoionisation with a subsequent Auger emission were taken into account. In the present paper we focus on photoionisation mechanisms, including shake-up with a subsequent satellite photoionisation (case a) and shake-off (cases b and c), which under certain conditions may produce multiple ionisations in the atom (case c). We investigate in detail what effects these processes may have on model compounds in order to assess radiation induced damage processes in biological samples. We calculate the different shake contributions for carbon, nitrogen, and oxygen atoms in a model peptide (see Fig. 1a). Shake effects for sulphur are estimated in separate calculations for the three most common chemical environments of sulphur in proteins (Cys, Met, Cys-S-S-Cys; Figs. 1b-d).

Results

Fig. 1 shows the model compounds used in the calculations. In a recent polymer study (Nakayama *et al.*, 1999), experimentally observed differences in intensity for ester and carbonyl oxygen main lines could be explained directly from consideration of the main line intensities, where the main line intensity was estimated from the overlap obtained from the $n = 0$ term in equation (1). This has the clear advantage for inherently large molecules, such as polymers or polypeptides, that main-line losses can be estimated without consideration of all the shake states to which intensity is distributed. For large molecules, the number of shake states goes beyond that which can readily be calculated at the configuration interaction level of theory. In a Gly-Gly -Gly tripeptide (Fig. 1a), the central glycine unit can be expected to display a shake spectrum which is similar to that of a residue within a longer polypeptide chain. To investigate effects arising from the truncations of the polypeptide chain, shake effects were calculated for all atoms in the model, including the atoms from the terminal carboxyl and amino functional groups.

There are six **carbon** atoms in the peptide model, showing very similar overlaps between the initial unionized state, and the core ionized ground state (cf. Table 1). This overlap is ~ 0.85 for all carbons, except for the carbon at the terminal carboxylic acid group, which has an overlap of ~ 0.87. The intensity loss from the main line is therefore close to ca. 30% for all carbon atoms in these models (cf. Table 1).

The **oxygen** shake contribution is generally the largest. In particular, the carbonyl oxygens display large shake effects, in agreement with the general behaviour for organic molecules. Here, strong shake is found for the two carbonyl oxygens in the peptide links, as well as for the double-bonded oxygen in the terminal carboxylate functional group. The overlap for these three oxygens is ~ 0.78 (cf. Table 1), corresponding to a

total shake intensity loss of ca. 40% (cf. Table 1). The terminal hydroxyl oxygen has a slightly larger overlap of ~ 0.82, corresponding to the main line intensity loss of ca. 33%.

There are two **nitrogens** in the peptide links, and one in a terminal amino group. The shake effect for all these atoms is similar: the overlap is ~ 0.90 (Table 1) and the main line loss is ca. 20% (cf. Table 1). Nitrogen shows the smallest shake contribution, and the results agree well with data on a polyimide polymer (Nakayama *et al.*, 1999), which also showed little nitrogen shake effect.

Sulphur shake effects have been calculated for three different common chemical environments (Fig. 1b-d and Table 2). The largest loss was found for a disulphide bridge (R-S-S-R, 25 %), and the smallest one for the thiol terminal group (R-SH, 21 %).

The initial models used for the sulphur calculations were smaller than the models used for the other atom types. This could, in principle, be a cause of concern, as these type of calculations are known to underestimate shake contributions in extended systems. In order to test the stability of the numerical results, additional calculations were performed on models with increasing chain lengths (CH_3-SH, CH_3-CH_2-SH, and $CH_3-CH_2-CH_2-SH$). The results indicated negligible drift (not shown).

Discussion

Radiation damage prevents the structural determination of single biomolecules and other non-repetitive structures (like cells) at high resolutions in classical electron or X-ray scattering experiments (Henderson, 1990; Henderson, 1995). Analysis of time-dependent components in damage formation suggests that the conventional damage barrier can be substantially extended at extreme dose rates and ultrashort exposure times (Neutze *et al.*, 2000; Hajdu, 2000; Hajdu *et al.*, 2000; Hajdu & Weckert, 2001; Ziaja *et al.*, 2001). A quantitative description of damage formation and a detailed analysis of the ionisation dynamics of the sample are crucial for planning experiments at future X-ray lasers. Results described in this paper give the first assessments of shake contributions to the ionisation of carbon, nitrogen, oxygen and sulphur atoms within the most common chemical environments in a protein molecule. In previous applications, the calculated shakeup intensities were usually a few percentage units too high, so we judge the present results to provide upper bounds to the total shake intensities.

Data for noble gases show that shake-up and shake-off effects were of comparable magnitude (Svensson *et al.*, 1988; Armen *et al.*, 1985; Wark *et al.*, 1991). Calculations

on carbon, oxygen and nitrogen by Mohammedein *et al.*, 1993 and El-shemi & Hassan, 1997 show that photoemission accompanied by double Auger decay has nearly zero probability for these light elements. Such triple ionisations can thus be neglected in the ionisation dynamics of C, O, N compounds. Sulphur, on the other hand, behaves differently, and the probability of multiple ionisation ($n > 2$) after the ejection of a K-shell photoelectron is larger. The average charge left in sulphur after a single K-shell photoionisation is estimated to be about 4 (Mohammedein *et al.*, 1993). For instance, for S: $P(2) \sim 5\%$, $P(4) \sim 34\%$, $P(6) \sim 3\%$ (Mohammedein *et al.*, 1993; El-shemi & Hassan, 1997). This implies that for sulphur, multiple ionisation mechanisms should be taken into consideration in order to obtain more accurate predictions for the ionisation dynamics of proteins.

The energy of the photoelectrons, which emerge from shake-up processes is similar to the energy of photoelectrons released during outer-shell ionisation (Nakayama *et al.*, 1999). In these calculations the deviation was within 10-40 eV. As a consequence, these electrons can leave a submicroscopic sample in the same manner as outer-shell photoelectrons. The inelastic mean free path of these electrons is of the order of a few hundred Ångstroms (Ashley, 1990 and references therein; Ziaja *et al.*, 2001), and thus the damage by the departing electrons will only have to be considered in large samples where such electrons may become trapped and deposit energy.

Low energy shake-off electrons will behave similarly to Auger electrons, and may cause secondary ionisation in the sample. A detailed analysis of secondary electron cascades elicited by low energy electrons can be found in Ziaja *et al.*, 2001.

Materials and methods

The average total shake contributions were estimated from quantum chemical (QC) calculations, using the semi-empirical INDO/S-CI program ZINDO, developed by Zerner and co-workers (Ridley & Zerner, 1976; Bacon & Zerner, 1979; Zerner *et al.*, 1980), with standard parameterisation. A single determinant description was used for the initial (unionized) state of the system, while the excited core-ionized states were obtained from Configuration-Interaction calculations with Single excitations (CIS). The equivalent core approximation was used to represent the core ionized species.

The calculations are based on the sudden approximation (Aoberg, 1967) valid for high energy photons, in which the intensity of a given shake line is proportional to the square of the overlap $\langle \Psi_0 | \Phi_n \rangle$:

$$P_n = |\langle \Psi_0 | \Phi_n \rangle|^2, \tag{1}$$

where Ψ_0 is the wave function of the N-1 remaining electrons in the neutral molecule, and Φ_n is the (N-1)-electron wave function of the nth state of the ionized system. These intensities were calculated with the SHAKEINT (Lunell, 1987; Lunell & Keane, 1988) program package. This method has been successfully applied to a wide range of systems, including organic molecules (c.f. Lunell *et al.*, 1978; Sjögren *et al.*, 1992), C60 (Enkvist *et al.*, 1995), polymers (PMDA-ODA polyimide) (Nakayama *et al.*, 1999), and adsorbates on metals (CO adsorbed on Cu(100) surfaces) (Persson *et al.*, 2000).

Acknowledgements

We thank Svante Svensson for illuminating discussions and David van der Spoel for excellent comments. This work was sponsored by the Swedish Natural Science Research Council (NFR), the Swedish Research Council for Engineering Sciences (TFR), the EU-BIOTECH Programme and in part by the Polish Committee for Scientific Research with grants Nos. 2 P03B 04718, 2 P03B 05119 to B. Z. A. S. was supported by a STINT distinguished guest professorship. B. Z. is supported by the Wenner-Gren Foundations.

References

[]

[Aoberg, 1967] Aoberg T. 1967. Multiple excitation of a many-electron system by photon and electron impact. *Ann. Acad. Sci. Fenn. Series A, VI. PHYSICA, Nr. 303* .

[Armen *et al.*, 1985] Armen, G.B., Aoberg, T., Kh Rezaul Karim, Levin, J.C., Crasemann, B., Brown, G.S., Hsiung Chen, M., Ice, G.E. 1985. Threshold double photoexcitation of argon with synchrotron radiation. *Phys. Rev. Lett.* **54**:182–185.

[Bacon & Zerner, 1979] Bacon, A.D., Zerner, M.C. 1979. An intermediate neglect of differential overlap theory for transition metal complexes: Fe, Co and Cu chlorides. *Theor. Chim. Acta* **53**:21–54.

[Dyson, 1973] Dyson, N.A. 1973. X-rays in atomic and nuclear physics. *Longman, London* .

[El-shemi & Hassan, 1997] El-shemi, A., Hassan, A.S. 1997. Ejection probabilities of electrons as result of reorganization effects of inner-shell ionized atoms. *Jpn. J. Appl. Phys., Part 1* **36(6A)**:3726–3729.

[Enkvist *et al.*, 1995] Enkvist, C., Lunell, S., Sjögren, B., Brühwiler, P.A., Svensson, S. 1995. The C_{1s} shakeup spectra of buckminsterfullerene, acenaphthylene, and naphtalene, studied by high resolution X-ray photoelectron spectroscopy and quantum mechanical calculations. *J. Chem. Phys.* **103**:6333–6342.

[Hajdu, 2000] Hajdu J. 2000. Single molecule X-ray diffraction. *Curr. Op. Struct. Biol.* **10**:569–573.

[Hajdu *et al.*, 2000] Hajdu, J., Hodgson, K., Miao, J., van der Spoel, D., Neutze, R., Robinson, C., Faigel, G., Jacobsen, C., Kirz, J., Sayre, D., Weckert, E., Materlik, G., Szöke, A. 2000. Structural studies on single particles and biomolecules. *LCLS: The First Experiments. SSRL, SLAC, Stanford, USA* .

[Hajdu & Weckert, 2001] Hajdu, J., Weckert, E. 2001. Life Sciences - Scientific applications of XFEL radiation. *TESLA, the Superconducting Electron-Positron Linear Collider with an integrated X-ray Laser Laboratory. Technical Design Report., DESY, ISBN 3-935702-00-0* **5**:150–168.

[Henderson, 1990] Henderson, R. 1990. Cryoprotection of protein crystals against radiation-damage in electron and X-ray diffraction. *Proc. R. Soc. Lond. Biol. Sci.* **241**:6–8.

[Henderson, 1995] Henderson, R. 1995. The potential and limitations of neutrons, electrons and X-rays for atomic resolution microscopy of unstained biological molecules. *Q. Rev. Biophys.* **28**:171–193.

[Hubbel *et al.*, 1980] Hubbel, J.H., Gimm, H.A., Overbo, I. 1980. Pair, triplet, and total atomic cross-sections (and mass attenuation coefficients) for 1MeV-100GeV photons in elements Z=1 to 100. *J. Phys. Chem. Ref. Data* **9**:1023–1045.

[Krause & Oliver, 1979] Krause, M.O., Oliver, J.H. 1979. Natural width of atomic K and L level Kα X-ray lines and several KLL Auger lines. *J. Phys. Chem. Ref. Data* **8**:329–338.

[Lunell, 1987] Lunell, S. 1987. *Technical Report No. 892, Dept. of Quantum Chemistry, Uppsala University* .

[Lunell & Keane, 1988] Lunell, S., Keane, M.P. 1988. *UUIP Report No. 1183, Institute of Physics, Uppsala University* .

[Lunell *et al.*, 1978] Lunell, S., Svensson, S., Malmqvist, P.A., Gelius, U., Siegbahn, K. 1978. A theoretical and experimental study of the carbon 1s shakeup structure of benzene. *Chem. Phys. Lett.* **54**:420–424.

[Mohammedein *et al.*, 1993] Mohammedein, A.M.M., Reiche, I., Zschornack, G. 1993. Z-dependences in the reorganization of inner-shell ionized atoms. *Radiat. Eff. Defect. S.* **126**:309–312.

[Nakayama *et al.*, 1999] Nakayama, Y., Persson, P., Lunell, S., Kowalczyk, S.P., Wannberg, B., Gelius, U. 1999. High resolution X-ray photoelectron spectroscopy and INDO/S-CI study of the core electron shakeup states of pyromellitic dianhydride-4,4'-oxydianiline polyimide. *J. Vac. Sci. Technol. A* **17**:2791–2799.

[Neutze *et al.*, 2000] Neutze, R., Wouts, R., van der Spoel, D., Weckert, E., Hajdu, J. 2000. Potential for biomolecular imaging with femtosecond X-ray pulses. *Nature* **406**:752–757.

[Persson *et al.*, 2000] Persson, P., Bustad, J., Zerner, M.C. 2000. INDO calculations of small copper clusters, and CO adsorbed on copper (100) surfaces. *J. Comp. Chem.* **21**:1221–1228.

[Ridley & Zerner, 1976] Ridley, J., Zerner, M.C. 1976. Triplet States via Intermediate Neglect of Differential Overlap: Benzene, Pyridine and the Diazines. *Theor. Chim. Acta* **42**:223–236.

[Siegbahn *et al.*, 1969] Siegbahn, K., Nordling, C., Johansson, G., Hedman, J., Heden, P.F., Hamrin, K., Gelius, U., Bergmark, T., Werme, L.O., Manne, R., Baer, Y. 1969. ESCA applied to free molecules. *North Holland, Amsterdam* .

[Sjögren *et al.*, 1992] Sjögren, B., Svensson, S., de Brito, A.N., Correia, N., Keane, M.P., Enkvist, C., Lunell, S. 1992. The C_{1s} core shake-up spectra of alkene molecules: an experimental and theoretical study. *J. Chem. Phys.* **96**:6389–6398.

[Svensson *et al.*, 1988] Svensson, S., Eriksson, B., Martensson, N., Wendin, G., Gelius, U. 1988. Electron shake-up and correlation satellites and continuum shake-off distributions in X-ray photoelectron-spectra of the rare-gas atoms. *J. Elec. Spec. Rel. Phenom.* **47**:327–384.

[Wark *et al.*, 1991] Wark, D.L., Bartlett, R., Bowles, T.J., Robertson, R.G.H., Sivia, D.S., Trela, W., Wilkerson, J.F., Brown, G.S., Crasemann, B., Sorensen, S.L., Schaphorst, S.J., Knapp, D.A., Henderson, J., Tulkki, J., Aoberg, T. 1991. Correspondence of electron spectra from photoionization and nuclear internal conversion. *Phys. Rev. Lett.* **67**:2291–2294.

[Wiik, 1997] Wiik B.H. 1997. The TESLA project: an accelerator facility for basic science. *Nucl. Inst. Meth. Phys. Res. B* **398**:1–8.

[Winick, 1995] Winick H. 1995. The linac coherent light source (LCLS): a fourth generation light source using the SLAC linac. *J. Elec. Spec. Rel. Phenom.* **75**:1–8.

[Zerner *et al.*, 1980] Zerner, M.C., Loew, G.H., Kirchner, R.F., Mueller-Westerhoff, V.T. 1980. An intermediate neglect of differential overlap technique for spectroscopy of transition-metal complexes. Ferrocene. *J. Am. Chem. Soc.* **102**:589–599.

[Ziaja *et al.*, 2001] Ziaja, B., van der Spoel, D., Szöke, A., Hajdu, J. 2001. Auger-electron cascades in diamond and amorphous carbon. *to be published in Phys. Rev. B* .

Table 1. *Overlaps* ($\langle\Psi_0|\Phi_0\rangle$) *and average total shake electron intensity* ($1 - |\langle\Psi_0|\Phi_0\rangle|^2$). *Values were calculated for all carbon, oxygen, and nitrogen atoms ocurring in the model polypeptide (Fig. 1a).*

Atom	Overlap	Average shake intensity
C1	0.853	27%
C2	0.858	
C3	0.850	
C4	0.857	
C5	0.853	
C6	0.869	
O1	0.776	38%
O2	0.773	
O3	0.785	
O4	0.823	
N1	0.902	19%
N2	0.896	
N3	0.897	

Table 2. *Total shake contributions* $(1 - |\langle\Psi_0|\Phi_0\rangle|^2)$ *calculated for sulphur in three different chemical environments.*

Molecule	Shake intensity
$CH_3CH_2 - SH$	21%
$CH_3CH_2 - S - CH_2CH_3$	22%
$CH_3CH_2 - S - S - CH_2CH_3$	25%
Average	23%

Figure 1. *The model polypeptide for calculating the shake contributions for carbon, nitrogen, and oxygen atoms (Fig. 1a). Shake effects for sulphur were estimated in separate calculations for the three most common chemical environments of sulphur in proteins (Cys, Met, Cys-S-S-Cys; Figs. 1b-d).*

Persson et al. Fig. 1

a Gly-Gly-Gly:



b Model for cysteine:



c Model for methionine:



d Model for a disulphide bridge: